

The High-Energy Spectra of Accreting Black Holes: Observational Evidence for Bulk-Motion Infall

Chris Shrader^{1,2} and Lev Titarchuk^{1,3}

Received _____; accepted _____

¹Laboratory for High-Energy Astrophysics, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA; shrader@grossc.gsfc.nasa.gov,titarchuk@lheavx.gsfc.nasa.gov

²Universities Space Research Association, Lanham MD

³George Mason University/Institute for Computational Sciences and Informatics, Fairfax VA

ABSTRACT

We discuss the emergent spectra from accreting black holes, considering in particular the case where the accretion is characterized by relativistic bulk motion. We suggest that such accretion is likely to occur in a wide variety of black hole environments, where the strong gravitational field is expected to dominate the pressure forces, and that this likely to lead to a characteristic high-energy spectroscopic signature; an extended power-law tail. It is in the high (soft) state that matter impinging upon the event horizon can be viewed directly, and the intrinsic power-law is seen. Certain types of Active Galactic Nuclei (AGN) may represent the extragalactic analog of the high-soft state accretion, which would further support our ideas, demonstrating the stability of the ($\alpha \sim 1.8$) power-law. This stability is due to the asymptotic independence of the spectral index on the mass accretion rate and its weak dependence on plasma temperatures. We have computed the expected spectral energy distribution for an accreting black hole binary in terms of our three model parameters: the disk color temperature, a geometric factor related to the illumination of the black hole site by the disk and a spectral index related to the efficiency of the bulk motion upscattering. We emphasize that this is a fully self-consistent approach, and is not to be confused with the more common phenomenological methods employing additive power law and black-body or multi-color disk. A test of the model is presented using observational data from the Compton Gamma Ray Observatory and the Rossi X-Ray Timing Explorer, covering $\simeq 2 - 200$ keV for two recent galactic black hole X-ray nova outbursts. The resulting model fits are encouraging and, along with some observational trends cited from the literature, they support our bulk-motion hypothesis.

Subject headings: accretion — black hole physics — — radiation mechanisms:
nonthermal — relativity — stars: individual (GRO J1655-40, GRS 1915+105)

1. INTRODUCTION

Do black holes interact with an accretion flow in such a way that a unique observational signature can be identified – that is one which is entirely distinct from those associated with other compact objects, based solely on the radiation observed at infinity? This is a crucial question confronting both theoretical and observational astrophysicists today; for recent reviews of the astrophysics of black holes [see e.g., Liang & Narayan (1997), Zhang et al. (1997a)]. Certainly a large body of evidence has been accumulated which supports the *existence* of black holes, the most convincing arguments being those invoking dynamical mass determinations [e.g. Orosz & Bailyn (1997)]. Other arguments have recently been advanced suggesting that X-ray nova flux histories demonstrate the existence of black-hole horizons [Narayan, Garcia & McClintock, (1997)], and in AGN, asymmetrical line features have identified and interpreted as originating in massive black hole environments [Tanaka et al. 1995, Fabian et al. (1995)]. Also, quasi-periodic oscillations (QPOs) have been attributed to dynamical time scales associated with the innermost stable orbits in black hole binaries [Morgan, Remillard & Greiner (1997)].

A distinct feature of black hole spacetime geometries, as opposed to those associated with other compact objects, is the presence of the event horizon. Near the horizon the strong gravitational field is expected to dominate the pressure forces and thus drive the accreting material into free fall. In contrast, for other compact objects the pressure forces are dominant near the surface and the free fall state is absent. Recently, Titarchuk and Zannias (1998) (hereafter TZ98) have developed the relativistic radiative transfer theory demonstrating that high-energy photons are produced by upscattering from the converging inflow within a few Schwarzschild radii. Only some fraction of the radiation emitted by the accretion disk illuminates the converging inflow site. It can be such a situation that the radiation density (or pressure) determined by the injected energy of those soft disk photons

and by the weak amplification they experience [Titarchuk, Mastichiadis & Kylafis, (1997); hereafter TMK97] is much smaller than the Eddington value. *We argue that then this difference is crucial and it results in a unique observational signature for accreting black holes.*

As explained above, this signature originates from upscattering of low energy photons by fast moving electrons with velocities, v , approaching the speed of light, c . A soft photon of energy E , in the process of multiple scattering off the electrons, gets substantially blue-shifted to energy

$$E' = E \frac{1 - (v/c) \cos \theta}{1 - (v/c) \cos \theta'} \quad (1)$$

due to Doppler effect provided at least one photon is scattered in the direction of electron motion (i.e. when $\cos \theta' \approx 1$). For example, in the first scattering event we assume the direction of incident photon, θ_1 , is nearly normal to the electron velocity, and the direction of the scattered photon is nearly aligned with the electron velocity. In the process the its outward propagation through the converging-inflow medium, the angle between the photon and electron velocity increases. Thus, in the second event the cosine angle, $\cos \theta_2$, tends to approach zero. The angle of outgoing photon, θ'_2 , has to be large enough, in order for the Doppler boosted photon to reach an observer.

Any system having a disk structure around a compact object is expected to have a source of low-energy photons. *The boosted photon component is seen as the extended power-law at energies much higher than the characteristic energy of the soft photons. And it is entirely independent of the initial spectral and spatial distributions of the low-energy photons.* The spectral index of the boosted photon distribution is determined only by the mass accretion rate and the plasma temperature of the bulk flow. *The presence of this high-energy power-law component is a generic feature of the model.* A key ingredient in support of our claim comes from the exact relativistic transfer calculations describing the

Compton scattering of the low-energy radiation field of the Maxwellian distribution of fast moving electrons (TZ98). It was proven mathematically that the power law is always present as a part of the black hole spectrum over a wide energy range, extending up to 500 keV. A turnover in the spectrum at about this energy, i.e. at $E \lesssim m_e c^2$, is a prediction of our model. Other extended power-law components, which may be related to the relativistic electron motion, e.g. in a jet, are not uniquely constrained to this energy band because they are not tied with the electron rest mass $m_e c^2$. The observations with CGRO/OSSE could in principle confirm or refute this prediction. In practice, the data thus far obtained are signal-to-noise limited and cannot address this issue in a definitive manner.

In this letter we extract the most important points regarding the radiative transfer and present observational evidence which supports the model.

2. Bulk-Motion Spectral Models

It has been shown elsewhere (TMK97, TZ98, Chakrabarti & Titarchuk (1995); hereafter CT95) that two effects, the bulk motion upscattering and the Compton (recoil) downscattering (herein BMC), compete forming the hard tail of the spectrum as an extended power law. The soft part of the spectrum comes from the disk photons seen directly and a subset of those photons which escape from the BMC atmosphere after undergoing a few scatterings but without any significant energy change. It has also been shown that without taking into account special and general relativistic effects, one is able to reproduce the main features of the full relativistic formalism: the overall spectral energy distribution and the dependence of the high-energy power on mass-accretion rate (TMK97; TZ98); also, refer to recent calculations by Laurent & Titarchuk (1998) (hereafter LT98). In the relativistic treatment, the Compton downscattering becomes less efficient at high energies, due to Klein-Nishina effects. Also, there is a possibility that the electron

distribution in the converging inflow can deviate from a Maxwellian, flattening at high velocities, since there is insufficient time for it to thermalize. The hard photon power-law thus extends to higher energies. At the same time however, the spectrum is steepened as a result of gravitational redshift effects.

We have assumed that there is an external illumination of the converging flow by the low-energy black body radiation of an accretion disk having a characteristic temperature T_c . Furthermore we have assumed that this illuminating radiation impinges on the BMC atmosphere with a certain geometry, which we have parameterized in terms of a "fraction" f . This fraction is really the first expansion coefficient of the spatial source photon distribution over the set of the eigenfunctions of the BMC formulation (TMK97, Eq. 30). As we mentioned above (§1) the spectral index is independent of the illumination fraction, f . It is clearly demonstrated in TMK97 (Figs 4).

We remind the reader that all reasonable theoretical spectra must exhibit a smooth transition from blackbody-like spectrum to a pure power-law, typically, at energies in the 5–12 keV range for stellar black holes.

In the soft state when the accretion rate is higher, the soft photons from the the Keplerian disk cool the hot region (Compton cloud) due to thermal Comptonization and free-free emission (CT95). The cooler converging inflow, as it rushes towards the black hole, scatter the soft-photons within the a radius, $r \sim \dot{m}r_s$ – some of the photons then undergo outward radiative diffusion. Here $\dot{m} = \dot{M}/\dot{M}_E$, r_s is Schwarzschild radius, $\dot{M}$ is the net accretion rate (including accretion from the disk plus any halo or other non-keplerian component), $\dot{M}_E \equiv L_E/c^2 = 4\pi GMm_p/\sigma_T c$ is the Eddington accretion rate, M is the mass of the central object, m_p is the proton mass and G is the gravitational constant. It transfers its momentum to the soft-photons to produce the power-law component extending to energies comparable to the kinetic energy of electrons in the converging inflow, i.e. of

order $m_e c^2$. On the other hand *in the hard state, the hot emission cloud covering the BMC zone prevents us from seeing the photons that are upscattered to subrelativistic energies within a few Schwarzschild radii.*

The luminosity of the upscattered component, the hard power law, has to be very small compared to the Eddington luminosity in order for the BMC model to be valid. The relative normalization of the soft component to the hard power-law is less important provided the inferred luminosity of the hard power-law remains consistent with the assumption of negligible radiation pressure near the black-hole horizon.

The BMC spectral model can be described as the sum of a thermal (disk) component and the convolution of some fraction of this component $g(E_0)$ with the upscattering Green's function $I(E, E_0)$ (TMK97, Eq. 30). The Green's function has the form of a broken power-law with spectral indices α and $\alpha + \zeta$ for high $E \geq E_0$ and low $E \leq E_0$ energy parts respectively,

$$F_\nu(E) = \int_0^\infty I(E, E_0) g(E_0) dE_0. \quad (2)$$

The above convolution is insensitive to the value of the Green's function spectral index $\alpha + \zeta$, which is always much greater than one. TZ98 presented rigorous proof that the hard power-law tail is a signature of a Schwarzschild black hole. Furthermore, the same statement is valid for the case of a rotating (Kerr) black hole, although a higher mass accretion rate is required to provide the same efficiency for the soft photon upscattering.

We note that the processes of absorption and emission (as free-free or synchrotron radiation) can be neglected provided the plasma temperature of the bulk flow is of order 1 keV or greater for characteristic number densities of order 10^{18} cm^{-3} and for magnetic field strengths in the proximity of the black hole of order $10^4 - 10^5$ gauss or less (CT95).

3. Application to Recent High Energy Observations

As a test of the model we have collected data resulting from high-energy observations covering recent activity periods in two galactic X-ray novae: GRO J1655–40 [Zhang et al. (1997b)] and GRS 1915+105 [Chaty et al. (1996)]. X-ray novae comprise perhaps the best test case of the methodology described here, since they are in low-mass binary systems – avoiding the added complications which may arise from the OB star winds in high-mass binary BHCs such as Cygnus X–1 – and because they become exceptionally bright in outburst exhibiting frequent and pronounced high-energy spectral state transitions [e.g. Chen et al. (1997), Ebisawa et al. (1994) Esin, McClintock & Narayan, (1997)]. Furthermore, as a group they comprise the most convincing galactic black-hole candidates.

We constructed composite high-energy spectra for GRO J1655–40 during an outburst in the spring of 1996 (Hynes et al. (1998)), covering the 2–200 keV spectral region and fit these data by the BMC model. This was accomplished using summed, standard mode (128 channel) data from the RXTE/PCA and the 16-channel BATSE/LAD earth-occultation data bracketting the pointed observations. In addition, there was substantial outburst activity in GRS 1915+105 during the latter part of 1996 [e.g. Bandyopadhyay et al. (1998)]. We utilized some of the available data from the same instruments for this event as well.

The BMC model described in section 2 was imported into the "XSPEC" software package which was used to perform all of the model fitting described here. Our resulting fits are shown in Figure 1.

For GRO J1655-40 we obtained a blackbody color temperature of $kT_c = 1.1 \pm 0.1$ keV for the soft photon source, a energy spectral index of $\alpha = 1.60 \pm 0.03$, and a geometric factor f , parameterizing the fraction of the total soft photon flux illuminating the BMC inflow atmosphere, of $f = 0.32 \pm 0.02$. The observed 2-100 keV flux was 5.7×10^{-8} ergs/cm²/s. We note that the corresponding luminosity in the hard power-law component is about

1.5% of L_E (with an assumption of the distance to the source, 3.2 kpc and the mass of the central object, 7 solar masses), which is consistent with our assumption of negligible radiation pressure near the event horizon. Similar results; $kT_c = 0.9 \pm 0.1$, $\alpha = 1.68 \pm 0.03$ and $f = 0.72 \pm 0.02$, were obtained for GRS 1915+105.

From the inferred 2-200 keV luminosity for GRO J1655-40, $\sim 5\%$ of L_E , we derive a mass accretion rate (in Eddington units) of order 1, bearing in mind the efficiency of gravitational to radiative energy conversion is of order 5% or less, e.g. Shakura & Sunyaev 1973 (hereafter SS73). This value of $\dot{M}$ is consistent with expected values within the BMC framework. This suggests that the line-of-sight column density of the BMC atmosphere is of order 10^{24} cm^{-2} (see, TMK97, Eq. 2). However, because the best-fitted color temperature is about 1 keV, (and this is a lower bound on the BMC plasma temperature) we conclude that the detected X-ray spectrum is not significantly modified by absorption (see also §2).

The temperatures we infer for these two sources are somewhat lower than values previously reported [e.g. Zhang et al. (1997b)], however this is to be expected. As noted, our procedure represents a fully self-consistent model deconvolution, whereas most previous approaches are phenomenological, i.e. power law plus black body or multicolor disk. Mathematically, one expects the power law component contribute significant soft-energy flux with the net effect of skewing the thermal residual to higher apparent temperatures. This will not occur with approach, as the hard power law turns over towards low energies (see Fig 3, TMK97). The inferred spectral indices also agree extremely well with our model predictions. In TZ98, calculations of the $\dot{m} - \alpha$ relationship were presented. For the low temperature limit, an asymptotic lower limit of $\alpha \simeq 1.8$ was calculated; for the higher BMC plasma temperatures (of order 10 keV) this limit is significantly lower, ~ 1 , and for mass accretion rates of $\dot{m} \sim 1$ (see below), α is precisely in the 1.5 – 1.8 range we find (LT98, Titarchuk 1998). Thus, we feel our observational test provides extremely encouraging

support of our methodology.

Using our inferred color temperature T_c and spectral index α , along with the measured flux normalization, we can estimate the mass accretion rate, black hole mass and source distance within the framework of standard accretion disk theory (e.g. SS73). This additionally requires certain assumptions regarding a "hardening factor" – the ratio of color temperature to the effective plasma temperature (Shimura & Takahara (1995), hereafter ShT95). In fact, the spectral index (TMK97, TZ98, LT98) depends on the mass accretion rate and the plasma temperature; the disk color temperature is $\propto (\dot{m}/m)^{1/4}$ (SS73), and the normalization is $\propto \dot{m}m/d^2$ (where d is the distance to the source).

An ideal test case is GRO J1655-40, since its mass and distance are known to a high degree of accuracy (relative to other BHCs). The distance we infer, 3.8 ± 1.4 kpc, is consistent (at the $1 - \sigma$ level) with previous determinations (Hjellming, & Rupen (1995)). Also, the black hole mass we calculate can be reconciled with determinations from dynamical studies (Orosz & Bailyn (1997)) provided we used the hardening factor 1.9 (ShT95). This assumed a mass accretion rate $\dot{m} = 3$, which was the value obtained from Monte Carlo simulations performed to calculate the spectral index dependence on the mass accretion rate and plasma temperature (LT98). Again, this is an encouraging result suggesting that with further refinement, one has a method of mass and distance determination independent of the conventional quiescent spectroscopic and photometric studies, which are not always plausible.

4. DISCUSSION and CONCLUSION

The successful application of our basic model to observational data and the inferred physical parameters of those systems in comparison to independent determinations is

encouraging. We postulate that (i) *The soft state detected in GRO J1655-40 and GRS 1915+105 represents a generic feature of accreting Galactic black holes. An extragalactic analog may now be evident in the Narrow Line Seyfert 1 (NLS1) galaxy population.* (ii) *The BMC spectra represent a characteristic signature of black hole horizons. The disk flux, of order 5% L_{edd} tends to cool the ambient environment and the generic hard power-law components is seen by the observer.* (iii) *Because this spectral feature is formed very close to the horizon, $(2-3R_s)$, the variability timescales of high-energy line and continuum radiation should be associated with the crossing time scale $t_{\text{cross}} \sim 10^{-5}M/M_{\odot}$ s.* (iv) *The variability seen in the soft component is not expected to be correlated with the hard component. It is related to the illumination geometry of a small area of the black hole horizon site, whereas the soft radiation seen by the observer directly emanates from a major fraction of the entire disk which comprises a much larger area.* (v) *QPOs emanating from the inner edge of the accretion disk should lead to a pronounced hard- X-ray variability signature, because the seed photons for the converging flow upscattering come from the same inner disk region.* (vi) *The appearance of an additional bump in the energy range 10-20 keV can be explained in terms of downscattering (reflection) effects (ST80) from the inner edge of the accretion disk.*

Our conclusions are consistent with various observations of Galactic and extragalactic black hole systems. For example, in NLS1s the X-ray power-law is significantly steeper and its normalization is more variable, with time scale of order 10^4 s, than in broad-line Seyfert 1 galaxies. This suggests the NLS1s may represent the extragalactic analog of the high-soft state (Pounds, Done & Osborne (1995), Brandt, Mathur & Elvis (1997), Comastri et al. (1998)). Several groups have independently reached similar conclusions (CT95, Pounds, Done & Osborne (1995)). We further note that the equivalent widths of Fe features detected tend to be large, in some cases ~ 500 eV (Comastri et al. (1998), hereafter C98).

It is worth noting that the detection of the strong hydrogen-like iron line is expected

if the source of hard energy photons (> 7 keV) is located inside the converging inflow region (and alternatively, line radiation could form in the cooler, Compton cloud ambient to the converging inflow region). It is easy to show that the ionization parameter $\xi = L/(r^2 n) \approx 10^5$ ergs cm s^{-1} is a typical value for the converging inflow and it is almost independent of the central object mass. In this case, only the hydrogen-like iron would be expected (Kallman and McCray 1982), which has been confirmed recently by BeppoSAX observations (C98).

Another supportive example is the black hole X-ray binary LMC X-3 which appears to always be in the high state. Its hard-tail component varies independently of the soft component [e.g., Ebisawa et al. (1993)].

Recent RXTE observations of Cyg X-1 during a state transition [Cui et al. (1997)] revealed a striking decrease in the soft-to-hard photon lag times as the source passes from the hard to soft state. This is very strong empirical evidence that the soft seed photons, which comprise a of fraction the disk thermal component, and the hard-X-ray power law emanate from a common compact region, again consistent with our model.

The detection of 67 Hz QPOs from GRS 1915+105 by RXTE was recently reported by Morgan, Remillard & Greiner (1997). It was clearly demonstrated that this feature is associated with the high energy component visible in the PCA. This can be explained in terms of a QPO in the inner edge of the disk or by g -mode disk oscillations occurring within the characteristic radius of $4 r_s$ [Titarchuk, Lapidus & Muslimov (1998)]. This should lead to variations in the hard spectral component since significant changes in the illumination geometry of the converging inflow site, can occur (TMK97, TZ98). Similar interpretation can be applied to the 300 Hz QPOs detected by RXTE in GROJ1655–40 [Remillard et al. (1997)].

In conclusion, we wish to emphasize once again that the observations presented

here, along with some observational trends presented by others in the literature, and the relativistic theory prompt us to claim that *we have identified a generic spectral signature black hole accretion*.

ACKNOWLEDGMENTS

We wish to acknowledge the anonymous referee for who made a number of useful comments on the initial draft, as well as Jean Swank and Menas Kafatos for discussion and useful suggestions. Portions of this work were supported by the Rossi X-Ray Timing Explorer and Compton Gamma Ray Observatory Guest Observer Programs. L.T. also would like to acknowledge support from NASA grant NAG5-4965.

REFERENCES

- Bandyopadhyay, R., Martini, P., Gerard, E. Charles, P.A., Wagner, R.M., Shrader, C., Shahbaz, T., & Mirabel, I.F. 1998, MNRAS, in press
- Brandt, W.N., Mathur, S. & Elvis, M. 1997, MNRAS, 285, 25
- Chakrabarti S.K. & Titarchuk, L. G. 1995, ApJ, 455, 623 (CT95)
- Chaty, S., Mirabel, I. F., Duc, P. A., Wink, J. E., Rodriguez, L. F., 1996, A&A, 310, 825
- Chen, W., Shrader, C.R., & Livio, M., 1997, ApJ, 491, 312
- Comastri, A., Fiore, F., et al. 1998, A&A, in press (C98)
- Cui, W., Zhang, S.N., Focke, W., & Swank, J.H., 1997, ApJ, 484, 383
- Ebisawa, K., et al, 1993, ApJ, 403, 684
- Ebisawa, K., et al, 1994, PASJ, 46, 375
- Esin, A.A., McClintock, J.E., & Narayan, R., 1997, ApJ, 489, 865
- Fabian, A. C., Nandra, K., Reynolds, C. S., Brandt, W. N., Otani, C., Tanaka, Y., Inoue, H., Iwasawa, K. 1995, MNRAS, 277, L1
- Hjellming, R.M., & Rupen, M.P., 1995, it Nature, 375, 464.
- Hynes, R.I., Haswell, C. A., Chen, W., & Shrader, C. R., 1998, MNRAS, in press
- Inoue, H. et al., 1994, IAUC 6063
- Kallman, T.R. & McCray, R. 1982, ApJS, 50, 263
- Laurent, Ph., & Titarchuk, L.G., 1998 in preparation (LT98)

- Liang, E., & Narayan, R., 1997, in *Proceedings of the Fourth Compton Symposium* (eds Dermer, C.D. Strickman, M.S., & Kurfess, J.D.), AIP CP-410
- Morgan, E.H., Remillard, R.A., & Greiner, J., 1997, ApJ, 482, 993
- Narayan, R., Garcia, M.R., & McClintock, J.E., 1997, ApJ, 478, L79
- Orosz, J., & Bailyn, C.D. 1997, ApJ, 477, 876
- Pounds, K.A., Done, C., & Osborne, J., 1995, MNRAS, 277, L5
- Remillard, R.A., Morgan, E.H., McClintock, J.E., Bailyn, C.D. Orosz, J.A., & Greiner, J., 1997, (astro-ph/9705064)
- Shakura, N.I., & Sunyaev, R.A., 1973, A&A, 24, 337 (SS73)
- Shimura, T., & Takahara, F., 1995, ApJ, 445, 780 (ShT95)
- Sunyaev, R.A., & Titarchuk, L.G. 1980, A&A, 86, 121 (ST80)
- Tanaka, Y., Nandra, K., Fabian, A. C., Inoue, H., Otani, C., Dotani, T., Hayashida, K., Iwasawa, K., Kii, T., Kunieda, H., Makino, F., Matsuoka, M., 1995, Nature 375, No.6533, 659
- Titarchuk, L., & Zannias. T., 1998, ApJ, 493, 863 (TZ98)
- Titarchuk, L. G., Mastichiadis, A., Kylafis, N. D. 1997, ApJ, 487, 834 (TMK97)
- Titarchuk, L.G., Lapidus, I.I. & Muslimov, A., 1998, ApJ, 499, (astro/ph-9712348)
- Titarchuk, L.G., 1998 in preparation
- Zhang, S.N., Mirabel, I.F., Harmon, B.A., Kroeger, R.A., Rodriguez, L.F., Hjellming, R.M., & Rupen, M.P., 1997a in *Proceedings of the Fourth Compton Symposium* (eds Dermer, C.D. Strickman, M.S., & Kurfess, J.D.), AIP CP-410

Zhang, S.N., et al, 1997b, ApJ, 479, 381

Figure 1. RXTE/PCA and BATSE/LAD spectrum for (a) Broad-band high-energy spectrum of GRO J1655-40 obtained on 20 June, 1996. The data are fitted to the bulk-motion accretion model. The resulting model parameters are as stated in the text. (b) Same instrumentation and methodology applied to November 1996 active state in GRS 1915+105



